

Isolation and identification macrocyclic spermine alkaloid(protoverbine) from verbascum speciosum

Majid Halimi^{a,*}, Malihe Nasrabadi^b

Department of Chemistry, Payame Noor University, P.O. BOX 19395-4697 Tehran, Iran

Received: 20 June 2016, Accepted: 3 April 2017, Published: 3 April 2017

Abstract

The 17-membered macrocyclic spermine alkaloid protoverbine (8-phenyl-1,5,9,13-tetra azacycloheptadecan-6-one) was isolated from the aerial part of *verbascum speciosum*. The natural polyamines are ubiquitous bases reported to have several important functions in animals, plants and microorganisms. They are involved in processes of transcription and translation in protein synthesis and influence growth and development (cell division, differentiation, embryogenesis, etc.). In plants, their involvement in organ development, flowering, fruit ripening, senescence and stress responses is reported. The structure of the protoverbine was established by the study of their spectral such as ¹HNMR, ¹³C-NMR, IR, ESI-MS analyses and chemical properties. To the best of our knowledge, this is the first report of protoverbine from this species.

Keywords: *Verbascum speciosum*, alkaloid, separation, protoverbine, macrocyclic spermine.

Introduction

Verbascum speciosum is a species of flowering plant in the figwort family known by the common name in Iran, Gole Mahur (Figure 1). It is native to Eastern Europe and Western Asia, known in many other regions as a roadside weed. It is a biennial herb forming a rosette of large leaves and an erect stem well exceeding one meter in maximum height. The leaves are 30 to 40 centimeters long and have smooth edges and pointed tips. The plant blooms in a large panicle with many branches lined with flowers. Each flower has a corolla measuring 2 to 3 centimeters wide with five yellow

petals. There are five stamens coated in long white hairs at the center. The fruit is a capsule up to 7 millimeters in length containing many seeds [1].

In the past few years, a number of macrocyclic alkaloids structurally derived from spermine have been found to occur in certain species of the families Acanthaceae, Ephedraceae, Fabaceae, Flacourtiaceae and Scrophulariaceae [2 -7].

To date, various alkaloids have been isolated from the aerial parts of some *verbascum* species, protoverbine, protomethine [8], verbaskine [9], anabesine, plantagonine, acetamide [10], verbacine, verballocine,

*Corresponding author: Majid Halimi

Tel: +98 (915) 3842199, Fax: +98 (583) 2297089

E-mail: majid_halimi@pnu.ac.ir

verbasitie, verbascenine, verbamedine, verdoline and verbasine
verballoscenine, verbaskin[11], [13].
verbametrine, isovebametrine [12],



Figure 1. *verbascum speciosum*

Since there are no reports on the isolated alkaloids of *Verbascum speciosum*, this paper has introduced macrocyclic spermine alkaloid, protoverbine, isolated for the first time from this species. Polyamines ubiquitously occur in all organisms and many fundamental cellular processes like replication, transcription, membrane stabilization, and modulation of enzyme activities

[14,15]. In plants, polyamines play an important role in development as well as responses to biotic and abiotic stress [16,17]. The polyamine spermine is a major natural intracellular compound capable of protecting DNA from free radical attack [18]. The structural elucidation of this alkaloid by means of spectroscopy analysis is described.




Figure 2. structural of protoverbine

Experimental

General

The NMR spectra were recorded on a FT-NMR Ultra shield BRUKER 400MHz spectrometer with TMS as the internal standard. IR spectra were obtained on an Avatar 370 FTIR

Thermo Nicolet and EI-MS spectra were obtained on a HP5973 Series Mass spectrometer. MPLC was performed on a Büchi 861equipment (Switzerland) comprised of a specification pump module C-601 equipped with a UV-Detector C-640 set

at a wavelength of 254 nm, a model 7725i sample injector equipped with a 5 μ L loop and a Büchi software package for data collection. Column that used in this assay was Glass Column C-690(4.6 \times 250mm) and fraction collector was C-660. The mobile phase was 10:1 (v/v) dichloromethane–methanol. All analyses were performed at a flow-rate of 0.5 mL/min with detection at 254nm. The mobile phase was filtered through a 0.45 μ m filter and degassed. Separations were performed at room temperature.

Plant material

Leaves of *Verbascum speciosum* were collected at the flowering stage from Birjand, Iran, in June 2010 and identified at the Research Center for Plant Sciences at Ferdowsi University of Mashhad, Iran. A voucher specimen has been deposited in the Environmental Department of Bojnourd Herbarium (EDBH:00102).




Figure 3. Scheme the extraction procedure

Extraction and isolation procedure

Alkaloids of the air-dried powdered aerial parts of *verbascum speciosum* were extracted following a classical protocol (Figure 2). Briefly, the plant material (808g) was macerated with EtOH (2.0 L) four times and filtered. The procedure was repeated until the negative test against Dragendroff's reagent. The EtOH extracts were concentrated to give 244.25g of crude extract (yield: 30.22%). The ethanol extract was acidified (pH = 2) with aqueous acetic acid 60% and the final volume was adjusted to 400 mL. The

aqueous acidic solution was then extracted with ethyl acetate (3 \times 200 mL) to remove neutral components.

After the removal of neutral components, the aqueous layer was basified with 25% NH₃ on the ice chest (pH 10-12) and extracted with CHCl₃(300 mL \times 10). CHCl₃ was removed under reduced pressure to give a crude alkaloidal fraction. The alkaloid extract (2g) which was subjected to a Medium-Pressure Liquid Chromatographies (MPLC) was performed using a Büchi 861 apparatus with RP18 and silica gel (70-230

mesh). For MPLC analysis, the crude extract was dissolved in CH₂Cl₂ / MeOH (10:1) and 10 μL were injected onto a column chromatography on silica gel (70-230 mesh), eluted with CH₂Cl₂-MeOH (10:1) isocratic to give one fraction.

Characteristics of protoverbine alkaloid

Solid (41mg), ¹H-NMR (400MHz, CDCl₃) δ:7.33-7.26 (m, 5H), 3.67-3.64 (m, 1H), 3.48-3.45(m, 1H), 3.11-3.08 (m, 2H), 2.90-2.51 (m, 11H), 1.74-1.70 (m, 2H), 1.57-1.53 (m, 2H), 1.44-1.41 (m, 4H).¹³C-NMR (100MHz, CDCl₃) δ:171.5, 128.4, 127.6, 126.8, 62.6, 47.4, 46.8, 46.1, 39.3, 39.1, 37.5, 31.5, 25.6, 25.2, FT-IR ν_{max} cm⁻¹: 3342, 1300, 1100, 1660 EIMS 70eV, m/z: 332.

Results and discussion

The alkaloids of *verbascum speciosum* were exhaustively extracted corresponding with Figure 2. The crude alkaloids were subjected to MPLC and eluted with solvent (CH₂Cl₂/MeOH) (10:1). A single MPLC separation step performed on the alkaloid extract gave one fraction. The structure elucidation of this compound was established by spectroscopic methods, including EI-MS, IR, ¹H-NMR and ¹³C-NMR experiments.

Macrocylic spermine alkaloid was isolated as a white to yellow needle crystal (m.p=187°C). The molecular formula C₁₉H₃₂N₄O was assigned to protoverbine based on mass spectrometry evidence EI-MS (positive ion) m/z 332[M⁺]. IR spectrum showed bands at 3342cm⁻¹ (N-H stretching vibration), 1300-1100 cm⁻¹ (C-N stretching) and 1660cm⁻¹ (carbonyl group C=O).

The ¹H-NMR spectrum of protoverbine showed ten signals, a multiplet at δ_H 7.33-7.26 in the

aromatic region, eight methylene multiplets (δ_H 1.41-3.48) and one methine multiplet (δ_H 3.65). The 14 carbon signals present in the ¹³C-NMR spectrum were assigned. The ¹³C-NMR spectrum revealed the presence of aromatic carbon at (δ_C 126.8-128.4), aliphatic carbon at (δ_C 25.2-62.6) and carbonyl carbon at (δ_C 171.5).

Conclusion

In summary, macrocylic spermine alkaloid was isolated from the leaves of *verbascum speciosum* with MPLC. As expected, protoverbine is the largest component of *verbascum speciosum* alkaloid isolated for the first time from this plant.

Acknowledgments

The authors are grateful to Dr. Nematollahi (Food and Drug Administration of The North Khorasan, Iran) for cooperating and to the Payame Noor University (PNU) for the financial support.

References

- [1] H. Vahedi, J. Lari, M. Halimi, M. Nasrabadi, A.R. Vahedi, *J ESSENT OIL BEAR PL.*, **2012**, *15*, 896-899.
- [2] S. Bienz, R. Detterbeck, C. Ensich, A. Guggisberg, U. Häusermann, C. Meisterhans, B. Wendt, C. Werner, M. Hesse, *Alkaloids Chem Biol.*, **2002**, *58*, 83-338.
- [3] H. Bosshardt, A. Guggisberg, S. Johne, H. Veith, M. Hesse, H. Schmid, *Pharm.Acta Helv.*, **1976**, *51*, 371.
- [4] P. Datwyler, H. Bosshardt, H. O. Bernhard, M. Hesse, S. Johne, *Helv.Chim.Acta.*, **1978**, *61*, 2646.
- [5] M. Tamada, K. Endo, H. Hikino, *Heterocyclics.*, **1979**, *12*, 783.
- [6] M. Tamada, K. Endo, H. Hikino, C. Kabuto, *Tetrahedron lett.*, **1979**, *20*, 873.

- [7] P. Datwyler, H. Bosshardt, H.O. Bernhard, M. Hesse, S. Johne, *Helv Chim. Acta.*, **1979**, 62, 2712.
- [8] A. Guggisberg K. Drandarov, M. Hesse, *Helv.Chim.Acta.*, **2000**, 83, 3035.
- [9] K. Ishihara , Y. Kuroki , N. Hanaki , S. Ohara and H. YamamotoZdena, *J. Am. Chem. Soc.*, **1996**, 118, 1569–1570.
- [10] R. Ziyaev, A. Abdusamatov, S.YU. Yunusov, *Chemistry of Natural Compounds.*, **1971**, 7, 842.
- [11] K. Drandarov, I.M.Hais, *Journal of chromatography A.*, **1996**, 724, 416-423.
- [12] K. Drandarov, A. Guggisberg, M. Hesse, *Helv.Chim.Acta.*, **1999**, 82, 229-237.
- [13] A. Guggisberg, K. Drandarov, M. Hesse, *Helv.Chim.Acta.*, **2000**, 83, 3035-3042.
- [14] R. Amooaghaie and S. Moghym, *African Journal of Biotechnology*, **2011**, 10, 9673-9679.
- [15] J. M.Tanguy, *Plant Growth Regulat.*, **2000**, 100, 675–688.
- [16] P.Roy, K. Niyogi, DN. Sen Gupta, B. Gosh, *Plant Sci.*, **2005**, 168,583-591.
- [17] S. Singh Gill and N. Tuteja, *Plant Signal Behav.*, **2010**, 5, 26–33.
- [18] S. Fujisawa, Y. Kadoma, *Anticancer Res.*, **2005**, 25, 965-9.